



INDICATIONS & USAGE

MONJUVI (tafasitamab-cxix), in combination with lenalidomide, is indicated for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL) not otherwise specified, including DLBCL arising from low grade lymphoma, and who are not eligible for autologous stem cell transplant (ASCT).

This indication is approved under accelerated approval based on overall response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

MANAGING TREATMENT WITH MONJUVI

MONJUVI is the only outpatient targeted immunotherapy for adult NTE patients with R/R DLBCL in 2L with 5-year data^{1,2*}

The 5-year analysis data from L-MIND have not been submitted to or reviewed by the FDA, and potential inclusion of these data in the final FDA-approved labeling has yet to be determined.



National Comprehensive Cancer Network[®] (NCCN[®]) recommends as an NCCN Category 2A, Preferred Treatment Option

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) recommend tafasitamab-cxix (MONJUVI) + lenalidomide as an NCCN Category 2A preferred second-line or subsequent treatment option (if not previously used) for certain[†] patients living with DLBCL.^{3†}

**An estimated
>4300
patients have received
MONJUVI in the US^{4§}**

[§]Since FDA approval in 2020.

[†]MONJUVI is a CD19-directed cytolytic monoclonal antibody.¹ *Category 2A treatment option for patients who relapsed in less than 12 months and are non-candidates for CAR T-cell therapy (excluding primary refractory disease); Category 2A treatment option for patients who relapsed after more than 12 months with no intention to proceed with transplant.³ †It is unclear if tafasitamab or loncastuximab tesirine or if any other CD19-directed therapy will have a negative impact on the efficacy of subsequent anti-CD19 CAR T-cell therapy.

NCCN makes no warranties of any kind whatsoever regarding their content, use, or application and disclaims any responsibility for their application or use in any way.

NTE=non-transplant eligible; R/R=relapsed/refractory; DLBCL=diffuse large B-cell lymphoma; 2L=second line.

IMPORTANT SAFETY INFORMATION

Contraindications

None.

Warnings and Precautions

Infusion-Related Reactions

MONJUVI (tafasitamab-cxix) can cause infusion-related reactions (IRRs).

In L-MIND, infusion-related reactions occurred in 6% of the 81 patients with DLBCL who received MONJUVI. Eighty percent of infusion-related reactions occurred during cycle 1 or 2. Signs and symptoms included fever,

chills, rash, flushing, dyspnea, and hypertension. These reactions were generally managed with temporary interruption of the infusion and/or with supportive medication.

Premedicate patients prior to starting MONJUVI infusion. Monitor patients frequently during infusion. Based on the severity of the infusion-related reaction, interrupt or discontinue MONJUVI. Institute appropriate medical management.

Please see the full [Prescribing Information](#) for more information about MONJUVI.

L-MIND: AN OPEN-LABEL, MULTICENTER, SINGLE-ARM, PHASE 2 STUDY WITH 5-YEAR FOLLOW-UP^{1,2,5}

L-MIND study design¹

- ▶ Efficacy and safety of MONJUVI in combination with lenalidomide followed by MONJUVI monotherapy were evaluated in adults with R/R DLBCL after 1 to 3 prior systemic DLBCL therapies, including a CD20-containing therapy
- ▶ Enrolled patients at the time of the trial were not eligible for or refused ASCT
- ▶ Efficacy was established in 71 patients with DLBCL, as assessed by an Independent Review Committee using the International Working Group Response Criteria (Cheson 2007)


Primary endpoint⁵

Best overall response rate (ORR) (CR + PR)

Select secondary endpoint⁵

Duration of response (DoR)

EACH CYCLE WAS 28 DAYS¹



Administer premedications, including acetaminophen, histamine H₁ receptor antagonists, histamine H₂ receptor antagonists, and/or glucocorticosteroids, 30 minutes to 2 hours prior to starting MONJUVI infusion to minimize infusion-related reactions.

R/R=relapsed/refractory; ASCT=autologous stem cell transplant; CR=complete response rate; PR=partial response rate; SD=stable disease.

IMPORTANT SAFETY INFORMATION

Warnings and Precautions (cont'd)

Myelosuppression

MONJUVI can cause serious or severe myelosuppression, including neutropenia, lymphopenia, thrombocytopenia, and anemia.

In L-MIND, among 81 patients with DLBCL who received MONJUVI, Grade 3 neutropenia was reported in 25%, Grade 3 thrombocytopenia in 12%, and Grade 3 anemia in 7%. Grade 4 neutropenia was reported in 25% and Grade 4 thrombocytopenia in 6%. Neutropenia led to treatment discontinuation in 3.7% of the patients. Febrile neutropenia occurred in 12%.

Monitor complete blood counts (CBCs) before each treatment cycle and throughout treatment. Monitor patients with neutropenia for signs of infection. Consider granulocyte colony-stimulating factor (G-CSF) administration. Withhold MONJUVI based on the severity of the adverse reaction. Refer to the lenalidomide prescribing information for dosage modifications.

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use

L-MIND EXAMINED PATIENTS WITH A BROAD RANGE OF CHARACTERISTICS^{1,4}

Select baseline characteristics (N=71) ^{1,4}		
Time between first DLBCL diagnosis and first documented relapse or progression	≤12 months	17 (23.9%)
	>12 months	53 (74.6%)
	Unknown	1 (1.4%)
IPI score at screening	0–2 (low and low-intermediate risk)	34 (47.9%)
	3–5 (intermediate-high and high risk)	37 (52.1%)
Refractory disease	Primary refractory disease*	14 (19.7%)
	Refractory to last prior therapy	32 (45%)
	Refractory to rituximab	30 (42%)
Prior CD20-containing therapy		100%
Median number of prior therapies		2
Prior lines of therapy	1	49%
	2 to 4	51%
Prior ASCT		9 (13%)
Median age (range)		71 years (41-86 years)
Race[†]	White	95%
	Asian	3%
Sex, male		55%
ECOG performance status	0	26 (36.6%)
	1	38 (53.5%)
	2	7 (9.9%)
Primary reasons patients were not candidates for ASCT	Age	47%
	Refractory to salvage chemotherapy	27%
	Comorbidities	13%
	Refusal of high-dose chemotherapy/ASCT	13%
Bulky disease[‡]		14 (20%)
Elevated LDH		40 (56.3%)

ECOG=Eastern Cooperative Oncology Group; IPI=International Prognostic Index; LDH=lactate dehydrogenase.

L-MIND included difficult to treat patients^{4,6,7}:

- ▶ **24%** of patients had ≤12 months between first diagnosis and first documented relapse or progression
- ▶ **52%** of patients had an IPI score of 3-5
- ▶ **20%** of patients had primary refractory disease

*Following a protocol amendment, primary refractory DLBCL was defined as no response to or progression/relapse during or within 6 months of frontline therapy. Prior to the amendment, only patients whose disease relapsed within 3 months were defined as primary refractory and excluded. Therefore, patients with disease that relapsed or progressed between 3 and 6 months of frontline therapy were recruited before the protocol amendment.⁸

[†]Race was collected in 92% of the 71 patients.¹

[‡]Data was collected in 70 patients.⁴

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
 tafasitamab-cxix | 200mg
 for injection, for intravenous use

HIGH ORR REACHED, WITH A MAJORITY OF RESPONDERS ACHIEVING CR


1-year primary analysis in patients with R/R DLBCL (N=71)^{1*}:

▶ **Best ORR: 55%** (n=39; 95% CI: 43%, 67%); **CR: 37%**; **PR: 18%**

MONJUVI, in combination with lenalidomide, was granted accelerated approval based on the 1-year primary analysis of the L-MIND study. The 5-year analysis data from L-MIND have not been submitted to or reviewed by the FDA, and potential inclusion of these data in the final FDA-approved labeling has yet to be determined.

5-year follow-up analysis^{9*}:

▶ **Best ORR: 54%** (n=38; 95% CI: 41%, 66%); **CR: 37%**; **PR: 17%**



Median time to response⁴

In the L-MIND study (n=71), the median time to response was **2.0 months**

1-year primary analysis in patients with R/R DLBCL (N=71)⁴

- ▶ Median time to **CR: 10.9 months** (n=25)
- ▶ Median time to **PR: 1.9 months** (n=26)

These analyses are exploratory in nature, and L-MIND was not designed or powered to evaluate and compare multiple subgroups. These results should be interpreted with caution given the small sample size and due to single-arm studies not adequately characterizing time-to-event endpoints, which may lead to estimates that are unstable.

¹Assessed by an Independent Review Committee.¹

3L=third line; CI=confidence interval.

The cutoff date for the primary analysis was November 30, 2018, and occurred after the last patient enrolled had completed 12 months of follow-up. The cutoff date for the 5-year follow-up analysis was November 14, 2022, and occurred after the last patient enrolled had completed 5 years of follow-up.^{2,4}

IMPORTANT SAFETY INFORMATION

Warnings and Precautions (cont'd)

Infections

Fatal and serious infections, including opportunistic infections, occurred in patients during treatment with MONJUVI and following the last dose.

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use


SUSTAINED REMISSION IN PATIENTS WITH R/R DLBCL^{1,4,9}

1-year primary analysis in patients with R/R DLBCL (N=71)^{1**}

► Median DoR 21.7 months (range: 0, 24)

MONJUVI, in combination with lenalidomide, was granted accelerated approval based on the 1-year primary analysis of the L-MIND study. The 5-year analysis data from L-MIND have not been submitted to or reviewed by the FDA, and potential inclusion of these data in the final FDA-approved labeling has yet to be determined.

5-year follow-up analysis: median DoR not reached (median follow-up 53.8 months [95% CI: 31.8-58.7])^{4,9††}



[†]Assessed by an Independent Review Committee.¹

^{††}Kaplan-Meier estimate.^{1,9}

^{†††}DoR rate at 5 years is a Kaplan-Meier estimate and should be interpreted with caution due to the small sample size and the number of censored patients.

The cutoff date for the primary analysis was November 30, 2018, and occurred after the last patient enrolled had completed 12 months of follow-up. The cutoff date for the 5-year follow-up analysis was November 14, 2022, and occurred after the last patient enrolled had completed 5 years of follow-up.^{2,4}

IMPORTANT SAFETY INFORMATION

Warnings and Precautions (cont'd)

Infections (cont'd)

In L-MIND, 73% of the 81 patients with DLBCL who received MONJUVI developed an infection. Grade 3 or higher infection occurred in 30%. Infection-related deaths occurred in 2.5% of patients, including a case of progressive multifocal leukoencephalopathy (PML). The most frequent Grade 3 or higher infection was pneumonia (7%). The most frequent infections of any grade were respiratory tract infections (51%, including pneumonias) and urinary tract infection (17%).

Monitor patients for signs and symptoms of infection and manage infections as appropriate. Consider infection prophylaxis per institutional guidelines. Consider treatment with subcutaneous or intravenous immunoglobulin (IVIG) as appropriate.

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use

MONJUVI: A WELL-ESTABLISHED SAFETY PROFILE WITH NO HOSPITALIZATION REQUIRED FOR ADMINISTRATION^{1,2}

- ▶ MONJUVI should be administered by a healthcare professional with immediate access to emergency equipment and appropriate medical support to manage infusion-related reactions. Monitor patients frequently during infusion. Hospitalization may be required to manage adverse reactions¹



- ▶ Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in clinical trials of a drug cannot be directly compared to rates in other clinical trials of another drug and may not reflect the rates observed in practice¹

- ▶ Serious adverse reactions occurred in 52% of patients who received MONJUVI¹
 - Serious adverse reactions in $\geq 6\%$ of patients included infections (26%), including pneumonia (7%) and febrile neutropenia (6%)
- ▶ Fatal adverse reactions occurred in 5% of patients who received MONJUVI, including cerebrovascular accident (1.2%), respiratory failure (1.2%), progressive multifocal leukoencephalopathy (1.2%), and sudden death (1.2%)¹
- ▶ Permanent discontinuation of MONJUVI or lenalidomide due to an adverse reaction occurred in 25% of patients and permanent discontinuation of MONJUVI due to an adverse reaction occurred in 15%¹
 - The most frequent adverse reactions which resulted in permanent discontinuation of MONJUVI were infections (5%), nervous system disorders (2.5%), and respiratory, thoracic, and mediastinal disorders (2.5%)
- ▶ Dosage interruptions of MONJUVI or lenalidomide due to an adverse reaction occurred in 69% of patients and dosage interruption of MONJUVI due to an adverse reaction occurred in 65%¹
 - The most frequent adverse reactions which required a dosage interruption of MONJUVI were blood and lymphatic system disorders (41%) and infections (27%)
- ▶ The most common adverse reactions ($\geq 20\%$) were neutropenia (51%), respiratory tract infection (51%), fatigue (38%), anemia (36%), diarrhea (36%), thrombocytopenia (31%), cough (26%), pyrexia (24%), peripheral edema (24%), and decreased appetite (22%)¹
- ▶ Clinically relevant adverse reactions in $<10\%$ of patients in L-MIND were¹:
 - Blood and lymphatic system disorders: lymphopenia (6%)
 - General disorders and administration site conditions: IRR (6%)
 - Infections: sepsis (4.9%)
 - Investigations: weight decreased (4.9%)
 - Musculoskeletal and connective tissue disorders: arthralgia (9%), pain in extremity (9%), musculoskeletal pain (2.5%)
 - Neoplasms: basal cell carcinoma (1.2%)
 - Nervous system disorders: headache (9%), paresthesia (7%), dysgeusia (6%)
 - Respiratory, thoracic, and mediastinal disorders: nasal congestion (4.9%), exacerbation of chronic obstructive pulmonary disease (1.2%)
 - Skin and subcutaneous tissue disorders: erythema (4.9%), alopecia (2.5%), hyperhidrosis (2.5%)

IRR=infusion-related reaction.

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use

L-MIND: ADVERSE REACTIONS IN 1-YEAR PRIMARY ANALYSIS¹

Adverse reactions (≥10%) in patients with R/R DLBCL who received MONJUVI in L-MIND

Adverse Reaction	MONJUVI in combination with lenalidomide (N=81)		
	All Grades (%)	Grade 3 or 4 (%)	
Blood and lymphatic system disorders	Neutropenia	51	49
	Anemia	36	7
	Thrombocytopenia	31	17
	Febrile neutropenia	12	12
Infections	Respiratory tract infection*	51	12
	Urinary tract infection [†]	17	4.9
General disorders and administration site conditions	Fatigue [‡]	38	3.7
	Pyrexia	24	1.2
	Peripheral edema	24	0
Gastrointestinal disorders	Diarrhea	36	1.2
	Constipation	17	0
	Abdominal pain [§]	15	1.2
	Nausea	15	0
	Vomiting	15	0
Respiratory, thoracic, and mediastinal disorders	Cough	26	1.2
	Dyspnea	12	1.2
Metabolism and nutrition disorders	Decreased appetite	22	0
	Hypokalemia	19	6
Musculoskeletal and connective tissue disorders	Back pain	19	2.5
	Muscle spasms	15	0
Skin and subcutaneous tissue disorders	Rash	15	2.5
	Pruritus	10	1.2

► No new safety signals were observed in the 5-year analysis²

*Respiratory tract infection includes: lower respiratory tract infection, upper respiratory tract infection, respiratory tract infection, bronchitis, pneumonia, nasopharyngitis, and related terms.¹

[†]Urinary tract infection includes: urinary tract infection, urinary tract infection bacterial, and related terms.¹

[‡]Fatigue includes asthenia and fatigue.¹

[§]Abdominal pain includes abdominal pain, abdominal pain lower, and abdominal pain upper.

^{||}Rash includes rash, rash maculopapular, rash pruritic, rash erythematous, and rash pustular.¹

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
 tafasitamab-cxix | 200mg
 for injection, for intravenous use

L-MIND: LABORATORY ABNORMALITIES IN 1-YEAR PRIMARY ANALYSIS¹

Select laboratory abnormalities ($\geq 20\%$) worsening from baseline in patients with R/R DLBCL who received MONJUVI in L-MIND

Laboratory Abnormality		MONJUVI in combination with lenalidomide*	
		All Grades (%)	Grade 3 or 4 (%)
Chemistry	Glucose increased	49	5
	Calcium decreased	47	1.4
	Gamma glutamyl transferase increased	34	5
	Albumin decreased	26	0
	Magnesium decreased	22	0
	Urate increased	20	7
	Phosphate decreased	20	5
	Creatinine increased	20	1.4
	Aspartate aminotransferase increased	20	0
Coagulation	Activated partial thromboplastin time increased	46	4.1

*The denominator used to calculate the rate was 74 based on the number of patients with a baseline value and at least one post-treatment value.

IMPORTANT SAFETY INFORMATION

Warnings and Precautions (cont'd)

Embryo-Fetal Toxicity

Based on its mechanism of action, MONJUVI may cause fetal B-cell depletion when administered to a pregnant woman. Advise pregnant women of the potential risk to a fetus. Advise women of reproductive potential to use effective contraception during treatment with MONJUVI and for 3 months after the last dose.


The combination of MONJUVI with lenalidomide is contraindicated in pregnant women because lenalidomide can cause birth defects and death of the unborn child. Refer to the lenalidomide prescribing information on use during pregnancy.

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
 tafasitamab-cxix | 200mg
 for injection, for intravenous use

SAFETY BY TREATMENT PHASE


HEMATOLOGIC TREATMENT-EMERGENT ADVERSE EVENTS (EXPOSURE-ADJUSTED)²



Please see the full [Prescribing Information](#) for more information about MONJUVI.

SAFETY BY TREATMENT PHASE

NON-HEMATOLOGIC TREATMENT-EMERGENT ADVERSE EVENTS (EXPOSURE-ADJUSTED)²



► Chart depicts treatment-emergent adverse events with at least 10 events in any treatment period

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
 tafasitamab-cxix | 200mg
 for injection, for intravenous use

DOSAGE AND ADMINISTRATION OF MONJUVI + LENALIDOMIDE

- ▶ MONJUVI should be administered by a healthcare professional with immediate access to emergency equipment and appropriate medical support to manage IRRs¹

The cycle length for MONJUVI is 28 days¹

▶ Cycle 1

DAYS	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
MONJUVI 12 mg/kg	■			■				■							■							■						
Lenalidomide 25 mg daily	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●							

▶ Cycles 2 and 3

DAYS	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
MONJUVI 12 mg/kg	■							■							■							■						
Lenalidomide 25 mg daily	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●						

▶ Cycles 4 to 12

DAYS	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
MONJUVI 12 mg/kg	■														■													
Lenalidomide 25 mg daily	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●						

▶ After 12 cycles, continue MONJUVI monotherapy until disease progression or unacceptable toxicity

DAYS	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
MONJUVI 12 mg/kg	■														■													

- ▶ Refer to the lenalidomide prescribing information for lenalidomide dosage recommendations, including for patients with renal insufficiency¹
- ▶ MONJUVI may be administered in a local office or clinic or at an outpatient center¹

IMPORTANT SAFETY INFORMATION

Adverse Reactions

Serious adverse reactions occurred in 52% of patients who received MONJUVI. Serious adverse reactions in ≥6% of patients included infections (26%), including pneumonia (7%) and febrile neutropenia (6%). Fatal adverse reactions occurred in 5% of patients who received MONJUVI, including cerebrovascular accident (1.2%), respiratory failure (1.2%), progressive multifocal leukoencephalopathy (1.2%), and sudden death (1.2%).


Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
 tafasitamab-cxix | 200mg
 for injection, for intravenous use


L-MIND: HEMATOLOGIC TOXICITIES MANAGEMENT AND DOSING MODIFICATIONS^{1,4,8}

Administer MONJUVI in combination with lenalidomide 25 mg orally for a maximum of 12 cycles, then continue MONJUVI as monotherapy until disease progression or unacceptable toxicity¹


Refer to the lenalidomide prescribing information for lenalidomide dosing recommendations, including for patients with renal insufficiency.¹



44% of patients in L-MIND received concomitant granulocyte-colony stimulating factor⁸



46% of patients (37/80) had at least one dose reduction of lenalidomide^{8*}



78% of patients (62/80) were able to receive a lenalidomide dose of ≥ 20 mg/day over the duration of their treatment^{4*}

- ▶ Permanent discontinuation of MONJUVI or lenalidomide due to an adverse reaction occurred in 25% of patients and permanent discontinuation of MONJUVI due to an adverse reaction occurred in 15%¹
- ▶ Dosage interruptions of MONJUVI or lenalidomide due to an adverse reaction occurred in 69% of patients and dosage interruption of MONJUVI due to an adverse reaction occurred in 65%¹

*81 patients were enrolled; 80 patients received MONJUVI plus lenalidomide and 1 received MONJUVI alone.⁸

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use

RECOMMENDED PREMEDICATIONS AND PROPHYLACTIC MEDICATION¹



Premedications

Administer premedications 30 minutes to 2 hours prior to starting MONJUVI infusion to minimize IRRs. Premedications may include acetaminophen, histamine H₁ receptor antagonists, histamine H₂ receptor antagonists, and/or glucocorticosteroids.

For patients not experiencing IRRs during the first 3 infusions, premedication is optional for subsequent infusions.

If a patient experiences an IRR, administer premedications before each subsequent infusion.

Thromboprophylaxis

Refer to the lenalidomide prescribing information for recommendations on prophylaxis for venous and arterial thrombotic events.

PREPARATION AND ADMINISTRATION OF MONJUVI

Reconstitute and dilute MONJUVI prior to infusion¹



RECONSTITUTION¹

1. Calculate the dose (mg) and determine the number of vials needed.
2. Reconstitute each 200 mg MONJUVI vial with 5 mL Sterile Water for Injection, USP with the stream directed toward the wall of each vial to obtain a final concentration of 40 mg/mL tafasitamab-cxix.
3. Gently swirl the vial(s) until completely dissolved. Do not shake or swirl vigorously. Complete dissolution may take up to 5 minutes.
4. Visually inspect the reconstituted solution for particulate matter or discoloration. The reconstituted solution should appear as a colorless to slightly yellow solution. Discard the vial(s) if the solution is cloudy, discolored, or contains visible particles.
5. Use the reconstituted MONJUVI solution immediately. If needed, store the reconstituted solution in the vial for a maximum of 12 hours either refrigerated at 36 °F to 46 °F (2 °C to 8 °C) or room temperature at 68 °F to 77 °F (20 °C to 25 °C) before dilution. Protect from light during storage.



DILUTION¹

1. Determine the volume (mL) of the 40 mg/mL reconstituted MONJUVI solution needed based on the required dose.
2. Remove a volume equal to the required MONJUVI solution from a 250 mL 0.9% Sodium Chloride Injection, USP infusion bag and discard it.
3. Withdraw the necessary amount of MONJUVI and slowly dilute in the infusion bag that contains the 0.9% Sodium Chloride Injection, USP to a final concentration of 2 mg/mL to 8 mg/mL. Discard any unused portion of MONJUVI remaining in the vial.
4. Gently mix the intravenous bag by slowly inverting the bag. *Do not shake*. Visually inspect the infusion bag with the diluted MONJUVI infusion solution for particulate matter and discoloration prior to administration.
5. If not used immediately, store the diluted MONJUVI infusion solution refrigerated for up to 18 hours at 36 °F to 46 °F (2 °C to 8 °C) and/or at room temperature for up to 12 hours at 68 °F to 77 °F (20 °C to 25 °C). The room temperature storage includes time for infusion. Protect from light during storage.

Do not shake or freeze the reconstituted or diluted infusion solutions.

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use

PREPARATION AND ADMINISTRATION OF MONJUVI (cont'd)



ADMINISTRATION^{1,8}

1. Administer MONJUVI as an intravenous infusion.¹
 - For the first infusion, use an infusion rate of 70 mL/h for the first 30 minutes, then, increase the rate so that the infusion is administered within 1.5 to 2.5 hours¹
 - In the L-MIND study, after the first 30 minutes, the rate of infusion was increased to 125 mL/h over a 2-hour period⁸
 - Administer all subsequent infusions within 1.5 to 2 hours¹
 - In the L-MIND study, vital signs were measured immediately prior to infusion, at 15 minutes (+/- 5 minutes), 30 minutes (+/- 10 minutes), every 60 minutes (+/- 15 minutes), and at the end of the infusion (+/- 20 minutes)⁸
2. Infuse the entire contents of the bag containing MONJUVI.¹
3. Do not co-administer other drugs through the same infusion line.¹
4. No incompatibilities have been observed between MONJUVI with infusion containers made of polypropylene (PP), polyvinylchloride (PVC), polyethylene (PE), polyethylene terephthalate (PET), or glass and infusion sets made of polyurethane (PUR) or PVC.¹

HOW MONJUVI IS SUPPLIED¹



- MONJUVI for injection is a sterile, preservative-free, white to slightly yellowish lyophilized powder for reconstitution, supplied as a 200-mg, single-dose vial
- Each 200-mg vial is individually packaged in a carton (NDC 73535-208-01 or NDC 50881-013-03)

STORAGE AND HANDLING OF MONJUVI¹



- Store refrigerated at 36 °F to 46 °F (2 °C to 8 °C) in the original carton to protect from light
- Do not shake
- Do not freeze

IMPORTANT SAFETY INFORMATION

Adverse Reactions (cont'd)

Permanent discontinuation of MONJUVI or lenalidomide due to an adverse reaction occurred in 25% of patients and permanent discontinuation of MONJUVI due to an adverse reaction occurred in 15%. The most frequent adverse reactions which resulted in permanent discontinuation of MONJUVI were infections (5%), nervous system disorders (2.5%), and respiratory, thoracic and mediastinal disorders (2.5%).

Please see the full [Prescribing Information](#) for more information about MONJUVI.







MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use

DOSAGE MODIFICATIONS FOR ADVERSE REACTIONS¹

► In the L-MIND study, IRRs occurred in 6% of the 81 patients. 80% of IRRs occurred during cycle 1 or 2

Management guidelines for IRRs and myelosuppression

Infusion-related reactions (IRRs)*

Severity	Dosage Modification
Grade 2 (moderate)	 Interrupt infusion immediately and manage signs and symptoms. <hr/>  Once signs and symptoms resolve or reduce to Grade 1, resume infusion at no more than 50% of the rate at which the reaction occurred. If the patient does not experience further reaction within 1 hour and vital signs are stable, the infusion rate may be increased every 30 minutes as tolerated to the rate at which the reaction occurred.
Grade 3 (severe)	 Interrupt infusion immediately and manage signs and symptoms. <hr/>  Once signs and symptoms resolve or reduce to Grade 1, resume infusion at no more than 25% of the rate at which the reaction occurred. If the patient does not experience further reaction within 1 hour and vital signs are stable, the infusion rate may be increased every 30 minutes as tolerated to a maximum of 50% of the rate at which the reaction occurred. <hr/>  If Grade 3 reaction returns, stop the infusion immediately and permanently discontinue MONJUVI.
Grade 4 (life-threatening)	 Stop the infusion immediately and permanently discontinue MONJUVI.

*Ensure premedications administered before subsequent infusions.

IMPORTANT SAFETY INFORMATION

Adverse Reactions (cont'd)





Dosage interruptions of MONJUVI or lenalidomide due to an adverse reaction occurred in 69% of patients and dosage interruption of MONJUVI due to an adverse reaction occurred in 65%. The most frequent adverse reactions which required a dosage interruption of MONJUVI were blood and lymphatic system disorders (41%) and infections (27%).

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use

DOSAGE MODIFICATIONS FOR ADVERSE REACTIONS (cont'd)¹

Management guidelines for IRRs and myelosuppression (cont'd)

Myelosuppression	
Severity	Dosage Modification
<p>Platelet count of 50,000/mcL or less</p>	<p> Withhold MONJUVI and lenalidomide and monitor CBC weekly until platelet count is 50,000/mcL or higher.</p> <hr/> <p> Resume MONJUVI at the same dose and lenalidomide at a reduced dose. Refer to lenalidomide prescribing information for dosage modifications.</p>
<p>Neutrophil count of 1,000/mcL or less for at least 7 days</p> <p>OR</p> <p>Neutrophil count of 1,000/mcL or less with an increase of body temperature to 100.4 °F (38 °C) or higher</p> <p>OR</p> <p>Neutrophil count less than 500/mcL</p>	<p> Withhold MONJUVI and lenalidomide and monitor CBC weekly until neutrophil count is 1,000/mcL or higher.</p> <hr/> <p> Resume MONJUVI at the same dose and lenalidomide at a reduced dose. Refer to the lenalidomide prescribing information for dosage modifications.</p>

CBC=complete blood count.

Refer to the lenalidomide prescribing information for lenalidomide dosage recommendations.

IMPORTANT SAFETY INFORMATION

Adverse Reactions (cont'd)

The most common adverse reactions (≥20%) were neutropenia (51%), respiratory tract infection (51%), fatigue (38%), anemia (36%), diarrhea (36%), thrombocytopenia (31%), cough (26%), pyrexia (24%), peripheral edema (24%), and decreased appetite (22%).

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
 tafasitamab-cxix | 200mg
 for injection, for intravenous use

COUNSELING YOUR PATIENTS¹

Advise the patient to read the FDA-approved patient labeling (Patient Information). Advise your patients to contact their healthcare provider if they experience signs and symptoms of:

▶ Infusion-related reactions

- Advise patients to contact their healthcare provider if they experience signs and symptoms of infusion-related reactions

▶ Myelosuppression

- Fever of 100.4 °F (38 °C) or greater, or signs or symptoms of bruising or bleeding should be reported immediately
- Advise patients of the need for periodic monitoring of blood counts

▶ Infections

- Fever of 100.4 °F (38 °C) or greater or signs or symptoms of infection should be reported immediately

▶ Embryo-fetal toxicity

- Advise pregnant women of the potential risk to a fetus. Women of reproductive potential should inform their healthcare provider of a known or suspected pregnancy
- Advise women of reproductive potential to use effective contraception during treatment with MONJUVI and for 3 months after the last dose
- Advise patients that lenalidomide has the potential to cause fetal harm and has specific requirements regarding contraception, pregnancy testing, blood and sperm donation, and transmission in sperm. Lenalidomide is only available through a REMS program

▶ Lactation

- Advise women not to breastfeed during treatment with MONJUVI and for 3 months after the last dose

REFERENCES: 1. MONJUVI Prescribing Information. Wilmington, DE: Incyte Corporation. 2. Duell J, Abrisqueta P, Andre M, et al. Tafasitamab for patients with relapsed or refractory diffuse large B-cell lymphoma: final 5-year efficacy and safety findings in the phase II L-MIND study. *Haematologica*. 2024;109(2):553-566. doi:10.3324/haematol.2023.283480 3. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) for B-Cell Lymphomas V.2.2025. © National Comprehensive Cancer Network, Inc. 2025. All rights reserved. Accessed February 12, 2025. To view the most recent and complete version of the guideline, go online to NCCN.org. 4. Data on File. Incyte Corporation. 5. ClinicalTrials.gov. A study to evaluate the safety and efficacy of lenalidomide with MOR00208 in patients with R-R DLBCL (L-MIND). <https://clinicaltrials.gov/study/NCT02399085>. Accessed February 25, 2025. 6. Crump M, Neelapu SS, Farooq U, et al. Outcomes in refractory diffuse large B-cell lymphoma: results from the international SCHOLAR-1 study. *Blood*. 2017;130(16):1800-1808. 7. McMillan AK, Phillips EH, Kirkwood AA, et al. Favourable outcomes for high-risk diffuse large B-cell lymphoma (IPI 3-5) treated with front-line R-CODOX-M/R-IVAC chemotherapy: results of a phase 2 UK NCRI trial. *Ann Oncol*. 2020;31(9):1251-1259. 8. Salles G, Duell J, González Barca E, et al. Tafasitamab plus lenalidomide in relapsed or refractory diffuse large B-cell lymphoma (L-MIND): a multicentre, prospective, single-arm, phase 2 study. *Lancet Oncol*. 2020;21(7):978-988. doi:10.1016/S1470-2045(20)30225-4 9. Duell J, Abrisqueta P, Andre M, et al. Tafasitamab for patients with relapsed or refractory diffuse large B-cell lymphoma: final 5-year efficacy and safety findings in the phase II L-MIND study. *Haematologica*. 2024;109(2) (Suppl):553-566. doi:10.3324/haematol.2023.283480

Please see the full [Prescribing Information](#) for more information about MONJUVI.

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use

MONJUVI IS THE ONLY OUTPATIENT TARGETED IMMUNOTHERAPY FOR ADULT NTE PATIENTS WITH R/R DLBCL IN 2L WITH 5-YEAR DATA^{1,2*}

INDICATIONS & USAGE

MONJUVI (tafasitamab-cxix), in combination with lenalidomide, is indicated for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL) not otherwise specified, including DLBCL arising from low grade lymphoma, and who are not eligible for autologous stem cell transplant (ASCT).

This indication is approved under accelerated approval based on overall response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).



We're Here to Support Your Eligible Patients During Treatment

Our mission is to help your patients start and stay on therapy by assisting with access and as-needed support.

Information and resources available through IncyteCARES for MONJUVI include:

- Benefits verification and prior authorization or appeal support
- Information about savings and financial assistance options[†]
- Practice resources and forms

For more information and resources for Healthcare Professionals, visit HCP.IncyteCARES/MONJUVI.

[†]Terms and conditions apply. Program terms may change at any time.



Questions?

Call IncyteCARES for MONJUVI
at 1-855-452-5234,
Monday through Friday,
8 AM–8 PM ET

RESOURCES FOR YOU AND YOUR PATIENTS

Create customized infusion schedules for your patients >

Download educational materials for healthcare professionals >

Find information and resources for your patients >

SELECT SAFETY INFORMATION

MONJUVI can cause serious adverse reactions including:

- **Infusion-Related Reactions:** MONJUVI (tafasitamab-cxix) can cause infusion-related reactions (IRRs). In L-MIND, infusion-related reactions occurred in 6% of the 81 patients with DLBCL who received MONJUVI. Eighty percent of infusion-related reactions occurred during cycle 1 or 2. Signs and symptoms included fever, chills, rash, flushing, dyspnea, and hypertension. These reactions were generally managed with temporary interruption of the infusion and/or with supportive medication.
- **Myelosuppression:** MONJUVI can cause serious or severe myelosuppression, including neutropenia, lymphopenia, thrombocytopenia, and anemia.
- **Infections:** Fatal and serious infections, including opportunistic infections, occurred in patients during treatment with MONJUVI and following the last dose.
- **Adverse Reactions:** Serious adverse reactions occurred in 52% of patients who received MONJUVI. Serious adverse reactions in $\geq 6\%$ of patients included infections (26%), including pneumonia (7%) and febrile neutropenia (6%). Fatal adverse reactions occurred in 5% of patients who received MONJUVI, including cerebrovascular accident (1.2%), respiratory failure (1.2%), progressive multifocal leukoencephalopathy (1.2%), and sudden death (1.2%).

Please see the full [Prescribing Information](#) for more information about MONJUVI.

*MONJUVI is a CD19-directed cytolytic monoclonal antibody.¹



MONJUVI and the MONJUVI logo are registered trademarks of Incyte.
Incyte and the Incyte logo are registered trademarks of Incyte.
All other trademarks are the property of their respective owners.
© 2025, Incyte. MAT-MON-00624 08/25