

MONJUVI

tafasitamab-cxix

rituximab &
lenalidomide

MONJUVI + R2—the first and only CD19- and CD20-targeted immunotherapy combination approved for **2L+ follicular lymphoma patients**¹

MONJUVI + R2 is a **12-cycle fixed duration treatment** that can be administered in a local outpatient setting **without hospitalization required**¹

inMIND was a Phase 3, double-blind, international, multicenter study of 548 adult patients with relapsed or refractory FL Grade 1, 2, or 3a after at least 1 systemic therapy, including an anti-CD20 antibody. Patients were randomized 1:1 to receive 12 cycles of MONJUVI + R2 or placebo + R2. The primary endpoint was investigator-assessed PFS using the Lugano criteria.^{1,2}

R2=rituximab and lenalidomide; 2L+=second-line plus; PFS=progression-free survival.

INDICATIONS & USAGE

MONJUVI (tafasitamab-cxix), in combination with lenalidomide and rituximab, is indicated for the treatment of adult patients with relapsed or refractory follicular lymphoma (FL).

Limitations of Use: MONJUVI is not indicated and is not recommended for the treatment of patients with relapsed or refractory marginal zone lymphoma outside of controlled clinical trials.

IMPORTANT SAFETY INFORMATION

Contraindications

None.

Warnings and Precautions

Infusion-Related Reactions

MONJUVI (tafasitamab-cxix) can cause infusion-related reactions (IRRs).

In inMIND, infusion-related reactions occurred in 16% of the 274 patients with FL who received MONJUVI in combination with lenalidomide and rituximab. Signs and symptoms included fever, chills, rash, flushing, dyspnea, and hypertension. These reactions were generally managed with temporary interruption of the infusion and/or with supportive medication.

Premedicate patients prior to starting MONJUVI infusion. Monitor patients frequently during infusion. Based on the severity of the infusion-related reaction, interrupt or discontinue MONJUVI. Institute appropriate medical management.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#).

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use

MONJUVI + R2 was studied in a 2L+ FL broad patient population that included tough-to-treat patients^{1,3-5}

In inMIND, tough-to-treat patients included **83% high tumor burden** (met at least 1 GELF criteria), **43% refractory to prior anti-CD20 therapy**, and **32% POD24**.^{1,6}

Baseline Patient Characteristics^{1,6,7}

Select Baseline Characteristic	MONJUVI + R2 (n=273)	Placebo + R2 (n=275)	
Median age, years (range)	64 (36, 88)	64 (31, 85)	
Sex, %	Male	55	54
	Female	45	46
Race, %*	White	80	80
	Asian	15	15
	Other races	1	2
FL grade at trial entry, % [†]	Grade 1	22	19
	Grade 2	52	55
	Grade 3a	25	26
At least 1 GELF criteria at baseline, %	81	84	
POD24, %	31	32	
Median number of prior lines of therapy (range)	1 (1, 7)	1 (1, 10)	
Refractory to most recent prior therapy, %	41	35	
Refractory to prior anti-CD20 therapy, %	43	42	

- Refractory lymphoma was defined as achieved less than PR to the last treatment or achieved a CR or PR that lasted <6 months⁷
- POD24 was defined as progression of disease within 24 months after initial diagnosis¹

FL=follicular lymphoma; GELF=Groupe d'Etude des Lymphomes Folliculaires; POD24=progression of disease within 24 months after initial diagnosis; PR=partial response; CR=complete response.

*Race was not reported in 4% (n=11) of patients in the MONJUVI + R2 arm and 4% (n=10) of patients in the placebo + R2 arm.⁷

[†]FL grade at trial entry was not reported for 3 patients in the MONJUVI + R2 arm and 1 patient in the placebo + R2 arm.⁷

IMPORTANT SAFETY INFORMATION

Warnings and Precautions (cont'd)

Myelosuppression

MONJUVI can cause serious or severe myelosuppression, including neutropenia, lymphopenia, thrombocytopenia, and anemia.

In inMIND, among 274 patients with FL who received MONJUVI in combination with lenalidomide and rituximab, new or worse Grade 3 or 4 cytopenias included decreased neutrophils in 48% (Grade 4, 19%), decreased lymphocytes in 22% (Grade 4, 1.8%), decreased hemoglobin in 9%, and decreased platelets in 8% (Grade 4, 4%). Febrile neutropenia occurred in 4.4%.

Monitor complete blood counts (CBCs) before each treatment cycle and throughout treatment. Monitor patients with neutropenia for signs of infection. Consider granulocyte colony-stimulating factor (G-CSF) administration. Withhold MONJUVI based on the severity of the adverse reaction. Refer to the lenalidomide prescribing information for dosage modifications.


Please see additional Important Safety Information throughout and the full [Prescribing Information](#).



59% risk reduction in disease progression or death with MONJUVI + R2 as assessed by IRC⁶

IRC-Assessed PFS (Secondary Endpoint)

Median PFS was not reached with MONJUVI + R2 vs 16 months with R2^{6*}



Number of patients at risk

MONJUVI + R2	273	260	246	210	200	162	113	98	72	58	28	20	12	3	2	0
R2	275	260	230	193	170	120	79	67	44	38	26	15	8	2	2	0

*PFS by IRC assessment was not formally tested for statistical significance.

Investigator-Assessed PFS (Primary Endpoint)

- Median PFS[†] was **22.4 months** (95% CI: 19.2, NE) with MONJUVI + R2 (n=273) vs **13.9 months** (95% CI: 11.5, 16.4) with R2 (n=275) (HR[‡]=0.43 [95% CI: 0.32, 0.58]; $P < 0.0001$) **after a median follow-up of 14.1 months[†]**
- 57% risk reduction in disease progression or death with MONJUVI + R2 vs R2**

- ORR was **84%** (95% CI: 79%, 88%) in the MONJUVI + R2 arm (n=228/273) vs **72%** (95% CI: 67%, 78%) in the R2 arm (n=199/275)[†]

CI=confidence interval; NR=not reached; NE=not evaluable; HR=hazard ratio.

[†]Kaplan-Meier estimates.¹⁶

[‡]Hazard ratio based on a stratified Cox proportional hazards model.¹⁶

IMPORTANT SAFETY INFORMATION

Warnings and Precautions (cont'd)

Infections

Fatal and serious infections, including opportunistic infections, occurred in patients during treatment with MONJUVI and following the last dose.


Among 274 patients with FL who received MONJUVI in combination with lenalidomide and rituximab in inMIND, Grade 3 or higher infections occurred in 24%, including fatal infections in 1.1% of patients.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#).

MONJUVI[®]
 tafasitamab-cxix | 200mg
 for injection, for intravenous use

Median time to next treatment not reached with MONJUVI + R2 vs 28.8 months with R2⁶

Exploratory Endpoint



Number of patients at risk

MONJUVI + R2	273	268	261	257	224	199	162	132	105	88	67	43	34	22	7	0	0
R2	275	268	248	233	199	166	124	101	78	62	43	30	23	13	5	2	0

Time to next treatment was defined as the time from randomization to start of next anti-lymphoma therapy for any reason or death due to any cause, whichever occurs first.⁷

Limitations of Analysis

The decision to initiate subsequent treatment was made by the treating physician and patient and is subject to variability based on investigator interpretation of patient and disease characteristics. The time to next treatment analysis is exploratory in nature and should be interpreted with caution.

*Kaplan-Meier estimates.⁶

[†]Estimated using a stratified Cox proportional hazards model.⁶

IMPORTANT SAFETY INFORMATION

Warnings and Precautions (cont'd)

Infections (cont'd)


The most frequent Grade ≥ 3 infections were respiratory tract infections (19%), including Grade 3 or higher pneumonia (14%) and COVID-19 infection (11%). Opportunistic infections of any grade occurred in 6% of patients including herpes simplex or zoster infection (5%), fungal pneumonia (1.1%, including *Pneumocystis jirovecii* pneumonia in 0.4%), and cytomegalovirus (CMV) reactivation (0.4%).

Monitor patients for signs and symptoms of infection and manage infections as appropriate. Consider infection prophylaxis per institutional guidelines. Consider treatment with subcutaneous or intravenous immunoglobulin (IVIG) as appropriate.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#).

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use

PFS benefit was consistent across patient subgroups with MONJUVI + R2^{6*}



- POD24 was defined as progression of disease within 24 months after initial diagnosis¹
- Refractoriness to prior anti-CD20 therapy was defined as not achieving a CR or PR to a prior regimen containing anti-CD20 mAb, or disease progression occurring during treatment with, or relapse <6 months after last dose of anti-CD20 mAb⁷

Limitations of analysis

The subgroup analysis is exploratory in nature, and inMIND was not designed or powered to evaluate and compare multiple subgroups. These results should be interpreted with caution.

*Analysis by investigator assessment.⁶

IMPORTANT SAFETY INFORMATION

Warnings and Precautions (cont'd)

Embryo-Fetal Toxicity

Based on its mechanism of action, MONJUVI may cause fetal B-cell depletion when administered to a pregnant woman. Advise pregnant women of the potential risk to a fetus. Advise women of reproductive potential to use effective contraception during treatment with MONJUVI and for 3 months after the last dose.

The combination of MONJUVI with lenalidomide and rituximab is contraindicated in pregnant women because lenalidomide can cause birth defects and death of the unborn child. Refer to the lenalidomide prescribing information on use during pregnancy.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#).



A comparable safety profile with MONJUVI + R2 vs R2 alone with no hospitalization required for administration^{1,6}

- MONJUVI should be administered with immediate access to emergency equipment and appropriate medical support to manage infusion-related reactions¹
- A similar rate of adverse events of any grade was reported for MONJUVI + R2 (99.3%) vs R2 alone (99.3%)⁶
- A similar rate of Grade 3/4 adverse events was reported for MONJUVI + R2 (71.2%) vs R2 alone (69.5%)⁷
- Adverse reactions that occurred more frequently (>5% difference) with MONJUVI + R2 vs R2 alone included COVID-19 infection, pneumonia, diarrhea, pruritus, fatigue, musculoskeletal pain, and mucositis¹

Adverse reactions (≥10%) in patients with relapsed or refractory FL who received MONJUVI in inMIND [*]		All Grades (%)		Grade 3 or 4 (%)	
		MONJUVI + R2 (n=274)	R2 (n=272)	MONJUVI + R2 (n=274)	R2 (n=272)
Infections	Respiratory tract infection	56	56	18	9
	COVID-19 infection [†]	34 [‡]	24 [‡]	10	2.9
	Pneumonia [‡]	18	11 [§]	14	7
	Upper respiratory tract infection [†]	17	22	1.1	0.4
Gastrointestinal disorders	Diarrhea	38	28	0.7	1.8
	Constipation	29	25	0.7	0
	Nausea	18	14	0.4	0.4
	Abdominal pain	13	18	0	2.2
Skin and subcutaneous tissue disorders	Rash [†]	37	33	3.6	1.5
	Pruritus	16	10	0.4	0
General disorders	Fatigue [†]	34	25	2.9	0.7
	Pyrexia	19	16	1.8	2.2
	Mucositis [†]	17	11	0.4	0
	Edema [†]	11	17	0.7	1.1
Musculoskeletal and connective tissue disorders	Musculoskeletal pain [†]	24	16	0.4	0.4
	Muscle contracture [†]	18	19	0	0
Respiratory, thoracic, and mediastinal disorders	Cough	21	19	0	0
Procedural complications	Infusion-related reaction [†]	16	16	0.7	0.4
Nervous system disorders	Peripheral neuropathy [†]	12	11	0	0.4
	Headache	10	7	0.4	0
Metabolism and nutrition disorders	Decreased appetite	10	9	0	0.7



The median number of treatment cycles completed with MONJUVI + R2 was 12 (range: 1-12).⁷



Discontinuation of MONJUVI or placebo due to an adverse reaction: MONJUVI + R2 (11%) vs placebo + R2 (7%).^{1,6}

*Table includes a combination of grouped and ungrouped terms. Adverse reactions were graded using NCI CTCAE version 5.0.¹

[†]Grouped term. See full Prescribing Information for other related terms.¹

[‡]Includes 2 fatal outcomes.¹

[§]Includes 3 fatal outcomes, including 2 reported under COVID-19 infection.¹

IMPORTANT SAFETY INFORMATION

Adverse Reactions

In the MONJUVI arm, serious adverse reactions occurred in 33% of patients, including serious infections in 24% of patients (including pneumonia and COVID-19 infection). Other serious adverse reactions in ≥ 2% of patients included renal insufficiency (3.3%), second primary malignancies (2.9%), and febrile neutropenia (2.6%). Fatal adverse reactions occurred in 1.5% of patients, including from COVID-19, sepsis, and adenocarcinoma.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#).

MONJUVI[®]
 tafasitamab-cxix | 200mg
 for injection, for intravenous use

Select laboratory abnormalities (>20%) worsening from baseline in patients with relapsed or refractory FL who received MONJUVI in inMIND¹

Laboratory Abnormality	All Grades (%)*		Grade 3 or 4 (%)*	
	MONJUVI + R2	R2	MONJUVI + R2	R2
Hematology				
Neutrophils decreased	75	71	48	44
Hemoglobin decreased	60	54	9	7
Lymphocytes decreased	57	51	22	19
Platelets decreased	40	43	8	9
Chemistry				
Alanine aminotransferase increased	47	42	1.5	1.5
Alkaline phosphatase increased	33	27	0	0
Creatinine increased	29	30	1.5	0.7
Aspartate aminotransferase increased	29	31	0	0.4
Glucose increased	28	28	0.4	1.1
Potassium decreased	24	24	3.3	3
Sodium decreased	24	22	1.5	0.4

*The denominator used to calculate the rate was 268-274 based on the number of patients with a baseline value and at least 1 post-treatment value.¹

IMPORTANT SAFETY INFORMATION

Adverse Reactions (cont'd)

Adverse reactions led to permanent discontinuation of MONJUVI in 11% of patients and dosage interruptions in 74%. The most frequent adverse reactions leading to dosage interruptions of MONJUVI were neutropenia (37% of all patients), COVID-19 (22%), pneumonia (11%), and infusion-related reaction (8%).

REFERENCES: **1.** MONJUVI Prescribing Information. Wilmington, DE: Incyte Corporation. **2.** A phase 3 study to assess efficacy and safety of tafasitamab plus lenalidomide and rituximab compared to placebo plus lenalidomide and rituximab in patients with relapsed/refractory (R/R) follicular lymphoma or marginal zone lymphoma (inMIND). ClinicalTrials.gov. Accessed April 4, 2025. <https://clinicaltrials.gov/study/NCT04680052> **3.** Link BK, Day B, Zhou X, et al. Second-line and subsequent therapy and outcomes for follicular lymphoma in the United States: data from the observational National LymphoCare study. *Br J Haematol*. 2019;184(4):660-663. doi:10.1111/bjh.15149 **4.** Casulo C, Dixon JG, Le-Rademacher J, et al. Validation of POD24 as a robust early clinical end point of poor survival in FL from 5225 patients on 13 clinical trials. *Blood*. 2022;139(11):1684-1693. doi:10.1182/blood.2020010263 **5.** Solal-Céligny P, Lepage E, Brousse N, et al. Doxorubicin-containing regimen with or without interferon alfa-2b for advanced follicular lymphomas: Final analysis of survival and toxicity in the groupe d'étude des lymphomes folliculaires 86 trial. *J Clin Oncol*. 1998;16(7):2332-2338. doi:10.1200/jco.1998.16.7.2332 **6.** Sehn L, Luminari S, Scholz C, et al. Tafasitamab plus lenalidomide and rituximab for relapsed or refractory follicular lymphoma: results from a phase 3 study (inMIND). Results presented at: 66th ASH Annual Meeting & Exposition; December 7-10, 2024; San Diego, CA. **7.** Data on file. Incyte Corporation. **8.** Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for B-Cell Lymphomas V.1.2026. © National Comprehensive Cancer Network, Inc. 2025. All rights reserved. Accessed December 23, 2025. To view the most recent and complete version of the guideline, go online to NCCN.org.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#).



Reach for MONJUVI for your 2L follicular lymphoma patients¹



Significant risk reduction in disease progression or death¹



In an exploratory analysis, median time to next treatment not reached with MONJUVI + R2 vs 28.8 months with R2^{6*}

- In the **MONJUVI + R2 arm, 78% did not receive subsequent treatment** at 2 years after randomization (estimated)⁷
- Time to next treatment analysis was not tested for statistical significance⁷



MONJUVI + R2 is a fixed duration treatment that can be administered in an outpatient setting without hospitalization required¹



A comparable safety profile with MONJUVI + R2 vs R2 alone^{1,6}

- Adverse reactions that occurred more frequently (>5% difference) with MONJUVI + R2 vs R2 alone included COVID-19 infection, pneumonia, diarrhea, pruritus, fatigue, musculoskeletal pain, and mucositis¹
- Serious adverse reactions occurred in 33% of patients who received MONJUVI + R2, including serious infections in 24% of patients (including pneumonia and COVID-19 infection). Other serious adverse reactions in $\geq 2\%$ of patients included renal insufficiency (3.3%), second primary malignancy (2.9%), and febrile neutropenia (2.6%)¹

*Time to next treatment was defined as the time from randomization to start of next anti-lymphoma therapy for any reason or death due to any cause, whichever occurs first.⁷



National Comprehensive Cancer Network® (NCCN®) recommends as an NCCN Category 1 Preferred Treatment Option⁸

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) recommend tafasitamab-cxix (MONJUVI) in combination with rituximab and lenalidomide as a **Category 1 preferred second-line** or subsequent therapy option (if not previously used) for FL after ≥ 1 prior systemic therapy, including an anti-CD20 mAb.

NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

For information about patient assistance, please visit [HCP.IncyteCARES/MONJUVI](https://www.incyte.com/hcp/monjuvi)

IMPORTANT SAFETY INFORMATION

Adverse Reactions (cont'd)

The most common adverse reactions ($\geq 20\%$) in patients receiving MONJUVI were respiratory tract infections (56%) (including COVID-19 infection and pneumonia), diarrhea (38%), rash (37%), fatigue (34%), constipation (29%), musculoskeletal pain (24%), and cough (21%). The most common Grade 3 or 4 laboratory abnormalities ($\geq 20\%$) were decreased neutrophils (48%) and decreased lymphocytes (22%).

Warnings and Precautions

Infusion-Related Reactions

MONJUVI can cause infusion-related reactions (IRRs).

In inMIND, infusion-related reactions occurred in 16% of the 274 patients with FL who received MONJUVI in combination with lenalidomide and rituximab. Signs and symptoms included fever, chills, rash, flushing, dyspnea, and hypertension. These reactions were generally managed with temporary interruption of the infusion and/or with supportive medication.

Premedicate patients prior to starting MONJUVI infusion. Monitor patients frequently during infusion. Based on the severity of the infusion-related reaction, interrupt or discontinue MONJUVI. Institute appropriate medical management.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#).



MONJUVI and the MONJUVI logo are registered trademarks of Incyte. Incyte and the Incyte logo are registered trademarks of Incyte. All other trademarks are the property of their respective owners. © 2026, Incyte. MAT-MON-00923 01/26

MONJUVI[®]
tafasitamab-cxix | 200mg
for injection, for intravenous use